

May 2008

FN/Food Preservation/2008-03

Principles of Boiling Water Canning

Kathy Riggs, FCS Agent *Brian A. Nummer*, Ph.D., Food Safety Specialist

The USDA Cooperative Extension system has been providing safe canning recommendations for over 100 years. It has been known since those early days that heat at the temperature of boiling water can destroy "almost" all microorganisms. Yeast, molds, and most bacteria are easily killed at at this temperature. The only microorganisms that survive are the bacteria that form protective spores, such as *Clostridium botulinum*.

High acid levels are used to prevent the sporeformers, especially, *Clostridium botulinum*, from growing. The measurement of how acid a food is called pH. When the pH is below 4.6 growth of sporeformers is inhibited.

Foods such as fruits, pickles, sauerkraut, jams, jellies, marmalades, and fruit butters/spreads fit into the *high acid* group since they have an acidity, or pH level, of 4.6 or lower. Most tomatoes and tomato products also fit into this category. However, some new varieties of tomatoes are borderline acid and the Cooperative Extension System now recommends adding a little extra acid as a safety measure.

**Tomato Acidification*: Current recommendations for acidification of whole, crushed, or juiced tomatoes, are to add 2 tablespoons of *bottled lemon juice* or ¹/₂ teaspoon of *citric acid* per quart of tomatoes. For pints, use 1 tablespoon bottled lemon juice or ¹/₄ teaspoon citric acid. Four (4) tablespoons of 5 percent acidity *vinegar* per quart may be used instead of lemon juice or citric acid.

Boiling Water Canner

Most boiling water canners are made of aluminum or porcelain-covered steel. They have fitted lids and removable racks that are either perforated or shaped wire racks. The canner must be deep enough so that at least one inch of briskly boiling water will be over the tops of jars during processing. A flat bottom must be used on an electric range. Either a flat or ridged bottom may be used on a gas burner. To ensure uniform processing of all jars with an electric range, the canner should be no more than 2 inches wider than the burner on any side.





Note: Some newer ceramic or glass top stoves prohibit the use of pots with a diameter larger than the burner and some prohibit the use of large stock pots due to the excessive weight. Check your manufacturer's recommendations before canning on these stoves.

Selecting the Correct Processing Time

To ensure the safety of food processed the boiling water bath method, use scientifically-tested recipes (dated 1988 or later) to prepare the produce and adjust the processing time for altitude using the chart below. Keep in mind that failing to properly **lengthen** processing times may result in short-term shelf-life or spoiled food due to the growth of mold, bacteria, and other microorganisms not detected by sight or smell. Food poisoning may result if the spoiled food is eaten.

Boiling Water Canner Altitude Chart

Normally, the processing times given for acid foods in recipes and canning guides are based on an altitude at or below 1,000 feet above sea level using the boilingwater method. At altitudes above 1,000 feet, adjust the processing time according to the chart below.

Altitude (feet)	Increase Processing Time
1,001 - 3,000	5 Minutes
3,001 - 6,000	10 Minutes
6,001 - 8,000	15 Minutes
8,001 - 10,000	20 Minutes

Steps for Successful Boiling Water Canning

1. Fill the canner halfway with water.

2. Preheat water to 140 degrees F. for *raw-packed* foods and to 180 degrees F. for *hot packed* foods. Raw- or cold-packed foods are placed directly into hot jars and covered with hot syrup; hot packed foods are partially cooked or heated through and placed hot into hot jars, and covered with hot syrup. See tested recipes for detailed instructions for various types of produce.

3. Load filled jars, fitted with lids, into the canner rack and use the handles to lower the rack into the water; or fill the canner, one jar at a time, with a jar lifter.

4. Add more boiling water, if needed, so the water level is at least 1 inch above jar tops.

5. Cover with the canner lid and turn heat to its highest position until water boils vigorously.

6. Set a timer for the number of minutes required for processing the food. (Check tested recipes for specific instructions for jams/jellies and pickles.)

7. Lower the heat setting to maintain a gentle boil throughout the process schedule.8. Add more boiling water, if needed, to keep the water level to above one inch of the jar lids.

9. When jars have been boiled for the recommended time, turn off the heat and remove the canner lid.

10. Using a jar lifter, remove the jars and place them on a protected surface, leaving at least 1-inch spaces

between the jars during cooling. Keep away from air drafts and let the jars cool at room temperature.

Sources: Adapted from information in the USDA *Complete Guide to Home Canning* and *National Center for Home Food Preservation.*



Photo: Canning workshop at USU May 2008.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions.

Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USUsponsored events and activities.

This publication is issued in furtherance of Cooperative Extension work. Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Noelle E. Cockett, Vice President for Extension and Agriculture, Utah State University.