

Figure 1. Daily maximum and minimum air and average soil temperatures at the line source pasture plots (Larry Johnson Farm) north of Randolph, UT, 1999.

The grass plots generally began to “green up” by mid April 1999. Fertilizer was applied on May 12 with a Gandy spreader. Phosphate (0-45-0) was applied at 162 lbs/acre perpendicular to the variety plots. The 1-N treatment of all grass varieties were fertilized with 103 lbs/acre nitrogen (34-0-0). Regar and NewHy 2-N plots received 206 lbs/acre and no nitrogen was added to the 0-N plots.

Neutron probe access tubes were installed in the various treatments as shown in Figure 2. Neutron probe readings in late July 1999 indicate root zone soil water depletion in the 24 to 30 inch depths for most varieties.

A serious ground squirrel infestation, combined with badger activity, influenced subplot harvest variability.

FIELD OBSERVATIONS

A visual appraisal rated all the varieties 5.5 or higher on vigor, stand density and color (scale 0-10) in early September 1997 after establishment. By early June 1998 the grass height was 18 inches or more for all plots except the Perennial Ryegrass and Matua Brome. The Matua did not survive the winter very well. Many of the Matua Brome plants came from seed in 1998 rather than as plants surviving the winter. By late June 1998 all the varieties, except the Perennial Rye and Tegmar Intermediate Wheatgrass, had fully emerged seed heads. Similar conditions were also observed by late June 1999.

SOIL WATER MEASUREMENTS

Neutron probe readings of soil water content at 6 inch depth increments were taken weekly beginning in early April 1999. Due to the abundant rain during April and May (3.7 inches) the first line source irrigation was May 25 with a second on June 11. As a result, Figure 3 shows that the soil water content was similar in both “wet” and “dry” areas of the plot until just prior to the first clipping on June 10, 1999.

The occurrence and depth of individual rain and irrigation events are also shown in Figure 3 for the period April 1 - mid October, 1999. The line source irrigation variation became evident in late May when soil water content in the “dry” (I2) treatment (Figure 4) dropped below 50% of available water in the 2 ½ foot root zone. Whereas, the soil water content in a “wet” (I5) treatment was maintained well above 50% depletion throughout the season (Figure 3).

IRRIGATION

The line source consisted of a single line of closely spaced sprinklers, 20 ft apart, from south to north through the center of the plots (Figure 2). The irrigation water was collected in cans spaced at 10 ft intervals across the center of each replication. Due to a cool, wet spring only one line source irrigation occurred (May 25) prior to the first harvest. Twelve irrigations applied approximately 35 inches of water 5 feet from the line varying down to about 1 inch at 55 feet from the center (see Figure 5). Irrigations occurred at approximately 7 day intervals beginning June 11. Irrigation amounts vary from 0.8 inches (I1 - West average of three catch cans) to 36 inches (I6 - East) as shown in Figure 5. The variation is an almost linear increase from dry to wet on each side of the plots. This variation is also reflected in root zone soil water content as shown in Figure 3 and Figure 4. An additional 8 inches of rainfall occurred from April 1 through September 15, 1999.

HARVEST SAMPLES 1999

The first clipping was taken on June 10, 1999. Plot subsamples (2 ft by 2 ft) were hand clipped in all variety and fertility treatments at 15 feet west and 15 feet east of the center making 72 samples total. A small plot harvester was used to clean up the plots. Water level treatment subplot samples were not taken due to the lack of an obvious line source effect resulting from the abundant rainfall. Yields varied from 0.3 ton/acre (Crown Blend Perennial Rye) to 2.6 ton/acre (Regar Meadow Brome - 2 x nitrogen) as given in Table 1. Nutrient analysis (NIR) for protein, ADF, NDF, RFV and TDN are given in Table 1. The protein content varied from about 14% (Regar 0-N) to 25% (BG-3 Perennial Rye and NewHy 2-N) and TDN ranged from about 56% (Regar 0-N) to 68% (BG-3). Both Perennial Ryegrasses (Crown Blend and BG-3) had relative feed values (RFV) over 160 while Regar 0-N was lowest at 105. Protein, TDN and RFV increased with additional amounts of nitrogen (Regar and NewHy).

YIELD

Dry matter yields (Table 1) at well watered plots were generally greater for the first cutting varying from 0.3 to 2.6 ton/acre than for the second cutting (1.1 ton/acre to 2.0 ton/acre), and greater for the second than the third (0.6 to 1.5 ton/acre).

The two exceptions were BG-3 and Crown Blend Perennial Ryegrasses, both of which had increases in yield from first to second cuttings. The yield of BG-3 also increased in the third cutting. This may be due to variability within the plot. These varieties may not have been growing vigorously early due to cool spring temperatures. There was also a significant difference in second and third cutting yields across the different irrigation levels, as expected.




Figure 3. Soil water content in 2 ½ foot root zone in “wet” (I5) line source irrigation water level treatment.




Figure 4. Soil water content in a 2 ½ foot root zone in “dry” (I2) line source irrigation water level treatments.

Table 1. Summary of Nutrient Analysis (NIR) & Dry Matter Yield for Irrigated Grass Pasture Plot near Randolph, Utah 1999 (*Note: Yields shown are from the 15 water levels*).

	Protein	ADF	NDF	RFV	TDN	Yield ton/acre	Protein lb/acre
First Cutting - 10 June 1999							
Regar Meadow Brome (2x Fertilizer)	22.12	24.02	54.87	119.16	63.68	2.62	1157
Regar Meadow Brome	19.69	26.71	56.83	111.65	61.07	2.51	987
Regar Meadow Brome (No Fertilizer)	13.68	29.37	58.56	105.26	55.79	1.88	514
NewHy (2x Fertilizer)	25.63	20.37	52.19	130.45	67.39	1.83	936
NewHy	24.41	21.75	52.66	127.26	66.06	1.62	789
NewHy (No Fertilizer)	16.96	25.37	58.08	110.82	59.42	1.12	380
Halmark Orchardgrass	21.73	23.98	53.46	122.40	63.40	1.80	780
Manchar Smooth Brome	23.39	23.86	53.83	121.85	64.68	2.11	985
Tegmar Intermediate Wheatgrass	22.59	22.45	51.33	129.52	64.50	1.57	709
BG-3 Perennial Ryegrass	25.43	18.10	44.63	160.12	67.90	0.44	226
Crown Blend Perennial Ryegrass	23.69	18.51	43.19	164.96	66.48	0.27	128
Matua Prairie Brome	20.43	24.94	52.19	124.69	62.14	1.41	577
Second Cutting - 15 July 1999							
Regar Meadow Brome (2x Fertilizer)	17.90	29.43	60.18	102.02	58.93	1.95	699
Regar Meadow Brome	15.20	31.98	61.79	96.62	56.16	1.77	538
Regar Meadow Brome (No Fertilizer)	12.06	34.15	61.30	94.75	53.16	1.36	328
NewHy (2x Fertilizer)	22.70	24.77	56.70	114.39	63.89	1.74	790
NewHy	18.96	27.56	59.38	106.13	60.27	1.67	632
NewHy (No Fertilizer)	15.11	31.18	59.76	100.94	56.32	1.07	322
Halmark Orchardgrass	17.25	29.77	60.31	101.42	58.34	1.70	587
Manchar Smooth Brome	19.74	27.11	56.05	112.89	60.98	1.05	416
Tegmar Intermediate Wheatgrass	15.94	30.91	61.80	97.91	57.02	2.02	645
BG-3 Perennial Ryegrass	19.00	26.73	51.62	124.11	60.55	1.21	460
Crown Blend Perennial Ryegrass	18.02	28.21	51.96	121.25	59.37	1.39	502
Matua Prairie Brome	15.60	30.66	55.70	109.38	56.84	1.35	422
Third Cutting - 14 September 1999							
Regar Meadow Brome (2x Fertilizer)	14.17	27.67	53.49	117.33	56.65	1.51	428
Regar Meadow Brome	12.79	28.48	54.70	113.58	55.37	0.94	239
Regar Meadow Brome (No Fertilizer)	11.95	29.50	52.17	118.04	54.45	0.68	163
NewHy (2x Fertilizer)	20.67	21.77	52.48	127.78	63.26	0.93	386
NewHy	17.24	23.04	49.50	133.40	60.31	0.61	209
NewHy (No Fertilizer)	15.78	23.30	50.27	131.90	59.15	0.90	284
Halmark Orchardgrass	12.85	25.72	50.49	126.89	56.24	1.64	422
Manchar Smooth Brome	17.58	22.62	44.99	147.60	60.69	0.54	190
Tegmar Intermediate Wheatgrass	15.23	25.33	49.51	130.13	58.14	0.68	207
BG-3 Perennial Ryegrass	13.29	24.58	42.91	151.31	56.90	1.33	355
Crown Blend Perennial Ryegrass	13.69	25.71	44.93	143.43	56.87	1.10	300
Matua Prairie Brome	13.83	25.08	41.17	158.37	57.16	0.85	236
Three Cutting Yield Weighted Average of Nutrients and Total Dry Matter Yield							
Regar Meadow Brome (2x Fertilizer)	18.79	26.66	56.23	113.20	60.41	6.08	2283
Regar Meadow Brome	16.93	28.82	58.13	106.89	58.38	5.21	1764
Regar Meadow Brome (No Fertilizer)	12.82	31.05	58.40	103.84	54.47	3.92	1005
NewHy (2x Fertilizer)	23.47	22.36	53.99	123.68	64.85	4.50	2112
NewHy	20.95	24.44	55.05	119.16	62.21	3.89	1630
NewHy (No Fertilizer)	15.98	26.77	56.38	113.55	58.30	3.09	986
Halmark Orchardgrass	17.41	26.45	54.78	116.88	59.33	5.14	1789
Manchar Smooth Brome	21.50	24.60	53.17	123.06	62.12	3.70	1591
Tegmar Intermediate Wheatgrass	18.27	26.91	56.00	114.64	59.89	4.27	1562
BG-3 Perennial Ryegrass	17.41	24.49	46.69	141.61	61.78	2.99	1041
Crown Blend Perennial Ryegrass	16.85	26.26	48.31	134.34	60.91	2.76	930
Matua Prairie Brome	17.07	27.11	50.90	126.90	58.71	3.62	1236

Note: Protein, %; ADF, Acid Detergent Fiber %; NDF, Neutral Detergent Fiber %; RFV, Relative Feed Value; TDN, Total Digestible Average of 3 replications for each (except 4 for NewHy & 2 for NewHy-No Fertilizer)

Total Irrigation Amounts Pasture Plots Randolph, UT 1999




Figure 5. Variation in irrigation application with the line source sprinkler May 25 through September 15, 1999, at the Randolph pasture plots.

Seasonal average protein varied from 12.8% to 23.5% and TDN varied from 54.5% to 65%, respectively, for Regar 0-N and NewHy 2-N. The Perennial Ryegrass had the highest RFV (134 and 142), whereas, Regar 0-N (RFV = 104), and Regar 1-N (RFV = 107) were the lowest.

Figure 6 shows that yields increased in proportion with the increase in the irrigation amounts. The average seasonal dry matter yield of the four varieties in the well watered plots (I6 East) was nearly double the yield of the dry plots (I1 West).

This relationship can also be seen in the effect of the increased nitrogen levels applied to the Regar Meadow Brome, with an obvious difference between the three treatments (Figure 7). The total dry matter yield production of 1-N levels show a 26% and 35% increase in yields compared to 0-N levels for NewHy and Regar, respectively (Table 1). The resulting increase in the 2-N treatment over the 1-N level is 16% and 14% for treatment. NewHy and Regar. Demonstrating a greater incremental increase in yield for the 1-N treatment.

Utah State University Extension is an affirmative action/equal employment opportunity employer and educational organization. We offer our programs to persons regardless of race, color, national origin, sex, religion, age or disability.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert L. Gilliland, Vice-President and Director, Cooperative Extension Service, Utah State University, Logan, Utah. (EP/06-2000/DF)

Estimated Yields of Selected Varieties
Pasture Plots near Randolph, UT 1999




Figure 6. Estimated season total dry matter yields of grass varieties at varying irrigation levels.

Estimated Yields - Regar Meadow Brome
With Differing Nitrogen Amounts




Figure 7. Estimated Season Total Dry Matter Yields of Regar Meadow Brome with varying Irrigation at three Nitrogen Levels.