



Opioid Overdose Response Training

Aaron Hunt, PhD & Gabriela Murza, MS, MCHES®
Extension Assistant Professors
Utah State University Extension
HEART Initiative



Purpose of Training

- This training focuses on overdose prevention and harm reduction best practices to help you know the signs and symptoms of an overdose, and feel confident in responding to an overdose.
- The training will last about 45 minutes and we will be collecting de-identified surveys from you to help us determine if this training is effective at educating you about overdose prevention.
- Before we can begin, we'll ask you to read and "sign" a consent letter and then take a pre-survey if you said "yes".
- You can still participate in the training if you indicate "no" on the consent form.





Outline

- Informed consent and pre-training survey
- Training
 - Overview of opioids and fentanyl
 - Identifying substance use
 - Signs of and responding to an opioid overdose
 - Legal considerations
 - Think-Pair-Share activity
 - Self-care and treatment resources
 - Post-training survey (if you consented)
- Next steps



Pre-Training Survey for SIE

- 1) Read the consent statement and indicate “yes” or “no” if you agree to complete the surveys
- 2) Answer questions to the best of your ability, even if you are not sure of your answer
- 3) Your answers will be kept confidential



https://usu.co1.qualtrics.com/jfe/form/SV_esvwe_xbtHMWVuUS



Pre-Training Survey for Students

- 1) Read the consent statement and indicate “yes” or “no” if you agree to complete the surveys
- 2) Answer questions to the best of your ability, even if you are not sure of your answer
- 3) Your answers will be kept confidential


https://usu.co1.qualtrics.com/jfe/form/SV_73cVjNQslhX5jYq



Opioid Epidemic In Utah


- 34-64 years old have highest risk
- 18-24 still have increased risk, especially among males

Poisoning: drug deaths by age and sex, Utah, 2019-2021




The Landscape in Different Parts of Utah

Poisoning: Drug deaths by Local Health District, Utah, 2018-2021



Highest Risk Occupations

Overdose Death Rate by Usual Industry Group 2020



([Billock et al., 2023](#))


Serve a Drink, Then Save a Life: This Is Restaurant Work During the Opioid Epidemic

As fentanyl overdoses in public spaces spike, the lifesaving drug Narcan is being stocked by more and more bars and restaurants.



Construction Industry Grapples With Its Top Killer: Drug Overdose

Construction workers are more likely to die of overdose than workers in any other occupation, forcing the industry to rethink its approach to safety.



Jump in elderly drug overdoses

A study in *JAMA Psychiatry* indicates that fatal drug overdoses in people 65 and older have quadrupled over the past 2 decades. Using CDC data to calculate the




Officials Warn of Fake Adderall Pills After Two College Students Die

Officials said the fake pills could contain fentanyl. Two Ohio State University students died this week in what the police said were apparent overdoses.



After several UNC-Chapel Hill students died from fentanyl, these students are handing out the antidote



DEA warns returning college students to be wary of potential fentanyl-laced pills

Federal officials are visiting Chicago-area campuses to spread the word about fake prescription pills that could be deadly.

What Are Opioids?

- Depressant - Help reduce pain by blocking pain receptors
- Opioids reduce consciousness and **decrease or stop breathing**
- The stronger an opioid, the more likely it is to cause an **overdose**



Oxycodone

OxyContin®, Oxecta®, Roxicodone®



Oxycodone/ Acetaminophen

Percocet®, Endocet®, Roxicet®



Hydrocodone/ Acetaminophen

Lortab®, Vicodin®, Lorcet®, Norco®



Codeine

Only available in generic form



Tramadol

ConZip®, Ultram®



Fentanyl

Actiq®, Duragesic®, Ventura®



Hydromorphone

Dilaudid®, Exalgo®



Meperidine

Demerol®



Methadone

Dolphin®, Methadone®



Morphine

MS-Contin®, Duramorph®, DepoDur®, Astramorph®



Heroin



DEA warns fentanyl most deadly drug in US and Utah history

By Heather Peterson and Curt Gresseth, KSL NewsRadio | Posted - March 28, 2024 at 8:42 a.m.



The DEA said it is on track to see double the amount of fentanyl in the state this year compared to last year. (Washington County Sheriff's Office)



Fentanyl

- A synthetic opioid that is 50 times more potent than heroin.
- Commonly comes in liquid or powder form.
- 2mg (fits on a pencil tip) causes overdose.
- Can't see, smell, or taste it.
- Unknown amount in and across substances.
- Tolerance is different for everyone.
- Commonly laced into street drugs
 - Not safe to experiment or use substances from drug dealers or online purchases.



Identifying Substance Use at Work, School, or Home

- Have you been in a situation where someone used opioids at your workplace, school, home, or someone's house (friend, family member, at a party)?
 - Where did they (or could they) use substances?
- Know how to access these locations
 - Knock loudly and ask for response
 - If unresponsive, use keys or tools to access a locked door
- Do you know what to do if someone experiences an opioid overdose? Do you know what the signs are?



Signs of an Opioid Overdose



Small, pinpoint pupils



Faint Heartbeat



Blue/purple fingernails & lips



Nonresponsive & can't be woken



Shallow or slow breathing



Gurgling, choking noise



Signs of Crystal Overdose



red face



very rapid or
irregular heartbeat



sweating heavily



rapid breathing, and/
or eye movement



aggression, anxiety,
extreme paranoia



hallucinations



fever



shaking or trembling
(jerky body movements)



stroke or heart attack

irregular breathing

pale, blue or clammy skin

vomiting

passed out, unconscious

seizures

confusion

ALCOHOL POISONING

save a life.

1. **+** KNOW THE FACTS
2. **#** RECOGNIZE AN EMERGENCY
3. **☎** CALL FOR HELP 518-580-5566

COCAINE OVERDOSES: WHAT YOU NEED TO KNOW

SIGNS AND SYMPTOMS OF A COCAINE OVERDOSE



RAPID
BREATHING
AND/OR
HEARTBEAT



MUSCLE
CRAMPING



WON'T
WAKE UP



CHEST
PAIN



FOAMING AT
THE MOUTH



SEIZURES OR
CONVULSIONS



DIZZINESS



DEHYDRATION



AGGRESSION,
ANXIETY
OR PARANOIA



SWEATING,
OFTEN WITH
CHILLS

EXTENSION.USU.EDU

Extension
UtahStateUniversity.



What is Naloxone/Narcan®?

- Naloxone – opioid overdose reversal medication
 - Narcan® - a brand of nasal spray
- Only works for opioid overdose; will not harm if opioids are not present
- No side effects, but watch for acute withdrawal
- Effects wear off in 30-90 minutes so still call 911
- May need to give more than one dose
- People often use multiple substances, so it is recommended to give Narcan® during overdose
- Available over the counter in most pharmacies, on store shelves, or free through public health programs



HOW DOES **NARCAN**[®] NASAL SPRAY WORK?



<https://narcan.com/resources>

EXTENSION.USU.EDU

Extension
UtahStateUniversity.



HOW TO USE



 **NARCAN**[®]
NASAL SPRAY




Steps to Responding to an Opioid Overdose

STEP 1: LAY



Check for breathing and responsiveness; lay on back.

STEP 2: SPRAY



Insert device into either nostril and press plunger firmly.

STEP 3: STAY




Stay with the individual and call 911 for help.

STEP 4: PROVIDE CHEST COMPRESSIONS




100 to 120 compressions per/minute ~ Stayin' Alive beat.

STEP 5: WATCH AND GIVE*



*If no response within 3 minutes, administer a second dose of naloxone.

STEP 6: PUT IN RECOVERY POSITION AND MONITOR RESPONSE



HAND SUPPORTS HEAD

KNEE STOPS ROLLING

References

Images: www.mass.gov/doc/mdph-oend-program-core-competencies/download

Opioid Overdose Response Information: www.narcan.com

Extension
UtahStateUniversity



HEALTH EXTENSION
Advocacy. Research. Teaching.

Will I Get in Trouble If I Call for Help?

“**Good Samaritan Laws**” in Utah protect people who call 911

- **Immunity from arrest for drug possession** when anyone calls 911 or seeks medical attention for an overdose
- BUT Utah does **not** protect people who report opioid overdoses from probation or parole violations



Think-Pair-Share

- **Harm Reduction:** A set of practical strategies and ideas aimed at reducing negative consequences associated with a behavior.
- Work with a partner to identify one way to promote harm reduction to a person who uses opioids.

COMMON EXAMPLES OF HARM REDUCTION




SUN
SCREEN



SEAT
BELTS



SPEED
LIMITS



BIRTH
CONTROL



CIGARETTE
FILTERS



Harm Reduction Strategies

- Carry Narcan or any naloxone product
- Don't mix opioids with alcohol or other substances
- Use less than usual
- Don't use street drugs or substances bought online
- Don't use alone in case of an emergency
 - If you're experimenting, ask for a buddy to be with you
 - If a friend is experimenting, offer to stay with them
- Store prescriptions safely – locked cabinet that's not easy to access
- Dispose of extras safely – don't keep them “just in case”. Can find a safe disposal site at <https://knowyourscript.org/> (scroll down the page).



Self-Care

- Responding to overdose emergencies are stressful situations
- If you need to speak with someone, there are local and national resources you can reach out to, including United Way's 211, and the Suicide and Crisis Lifeline at 988.



Treatment

- There are effective treatments for opioid use disorder.
 - Medications: Suboxone, methadone
 - Counseling/Therapy: Cognitive behavioral therapy and others



Please use this tool to find local substance use treatment providers



Call 211 or use link to be connected to local healthcare and social service resources



Provides recovery support resources for people and families recovering from substance use



Post-Training Survey for SIE

- Only take if you consented (indicated “yes” in the pre-survey)
- Answer the questions to the best of your ability
- Let us know if you have any questions

https://usu.co1.qualtrics.com/jfe/form/SV_42bMhd3KqK5qLXg



Post-Training Survey for Students

- Only take if you consented (indicated “yes” in the pre-survey)
- Answer the questions to the best of your ability
- Let us know if you have any questions

https://usu.co1.qualtrics.com/jfe/form/SV_3gZHPxPdkrSIR6S





Next Steps

EXTENSION.USU.EDU

Extension
UtahStateUniversity.




Overdose Prevention Resources

- Overdose response kits are given to organizations after a group training. It's kept onsite so anyone can respond to an overdose.
- Narcan® kits and wallet cards are given to attendees who complete a training.
 - In-person workshop - get it at the end.
 - Virtual workshop - will receive an email with information about pick-up locations.



Graphic Novel

- Tells a story of a server responding to an opioid overdose at work.
- Modified an existing graphic novel from a similar program in Atlanta, GA, with permission.
- Available at public locations and/or in a workshop.



Questions and Contact Information

- Dr. Aaron Hunt – USU Extension
 - Logan campus/statewide
 - Aaron.Hunt@usu.edu
 - 605-530-6900
- Gabriela Murza – USU Extension
 - Utah County
 - Gabriela.Murza@usu.edu
 - 385-268-6543



USU Extension - HEART



usuextension_heart



@usuHEART



USU HEART



<https://extension.usu.edu/heart>

References

- Algren D, Monteilh C, Rubin C, et al. Fentanyl-associated fatalities among illicit drug users in Wayne County, Michigan (July 2005-May 2006). *Journal Of Medical Toxicology: Official Journal of the American College Of Medical Toxicology* [serial online]. March 2013; 9(1):106-115.
- Centers for Disease Control and Prevention (CDC). [Vital signs: overdoses of prescription opioid pain relievers—United States, 1999–2008](#). *MMWR MorbMortal Wkly Rep*. 2011 Nov 4; 60(43):1487-1492.
- Chandler, R. K., Fletcher, B. W., & Volkow, N. D. (2009). Treating Drug Abuse and Addiction in the Criminal Justice System: Improving Public Health and Safety. *JAMA*, 301(2), 183–190. <https://doi.org/10.1001/jama.2008.976>
- Drug Enforcement Administration. 2019 National Drug Threat Assessment. Drug Enforcement Administration Strategic Intelligence Section, U.S. Department of Justice. Published December 2019. Accessed March 17, 2020 from https://www.dea.gov/sites/default/files/2020-01/2019-NDTA-final-01-14-2020_Low_Web-DIR-007-20_2019.pdf
- Doe-Simkins, M., Quinn, E., Xuan, Z., Sorensen-Alawad, A., Hackman, H., Ozonoff, A., & Walley, A. Y. (2014). Overdose rescues by trained and untrained participants and change in opioid use among substance-using participants in overdose education and naloxone distribution programs: a retrospective cohort study. *BMC Public Health*, 14, 297. <https://doi.org/10.1186/1471-2458-14-297>
- Ersche, K. D., Gillan, C. M., Jones, P. S., Williams, G. B., Ward, L. H. E., Luijten, M., ... Robbins, T. W. (2016). Carrots and sticks fail to change behavior in cocaine addiction. *Science*, 352(6292), 1468–1471. <https://doi.org/10.1126/science.aaf3700>
- Gladden RM, Martinez P, Seth P. [Fentanyl law enforcement submissions and increases in synthetic opioid-involved overdose deaths—27 states, 2013–2014](#). *MMWR MorbMortal Wkly Rep*. 2016; 65:837–43.



References

- Gladden M, O'Donnell J, Mattson C, Seth P. [Changes in Opioid-Involved Overdose Deaths by Opioid Type and Presence of Benzodiazepines, Cocaine, and Methamphetamine – 25 States, July-December 2017 to January-June 2018](#). MMWR Morb Mortal Wkly Rep 2019;68(34);737-744.
- Gottfredson, D. C., Najaka, S. S., & Kearley, B. (2003). Effectiveness of Drug Treatment Courts: Evidence from a Randomized Trial*. Criminology & Public Policy, 2(2), 171–196. <https://doi.org/10.1111/j.1745-9133.2003.tb00117.x>
- Jones, J. D., Campbell, A., Metz, V. E., & Comer, S. D. (2017). No evidence of compensatory drug use risk behavior among heroin users after receiving take-home naloxone. Addictive Behaviors, 71, 104–106. <https://doi.org/10.1016/j.addbeh.2017.03.008>
- Kariisa M, Scholl L, Wilson N, Seth P, Hoots B. [Drug Overdose Deaths involving Cocaine and Psychostimulants with Abuse Potential – United States, 2003-2017](#). MMWR Morb Mortal Wkly Rep 2019;68(17);41-43.
- Katzman, J. G., Takeda, M. Y., Bhatt, S. R., Moya Balasch, M., Greenberg, N., & Yonas, H. (2018). An Innovative Model for Naloxone Use Within an OTP Setting: A Prospective Cohort Study. Journal of Addiction Medicine, 12(2), 113. <https://doi.org/10.1097/ADM.0000000000000374>
- O'Donnell JK, Gladden RM, Seth P. [Trends in deaths involving heroin and synthetic opioids excluding methadone, and law enforcement drug product reports, by census region—United States, 2006–2015](#). MMWR MorbMortal Wkly Rep. 2017; 66:897–903.
- O'Donnell JK, Halpin J, Mattson CL, Goldberger BA, Gladden RM. [Deaths involving fentanyl, fentanyl analogs, and U-47700—10 states, July–December 2016](#). MMWR Morb Mortal Wkly Rep. 2017; 66:1197–202.



References

- Rees, D. I., Sabia, J. J., Argys, L. M., Latshaw, J., & Dave, D. (2017). With a Little Help from My Friends: The Effects of Naloxone Access and Good Samaritan Laws on Opioid-Related Deaths (Working Paper No. 23171). <https://doi.org/10.3386/w23171>
- Rudd RA, Paulozzi LJ, Bauer MJ, Burleson RW, Carlson RE, Dao D, Davis JW, Dudek J, Eichler BA, Fernandes JC, Fondario A. [Increases in heroin overdose deaths—28 states, 2010 to 2012](#). MMWR MorbMortal Wkly Rep. 2014 Oct 3; 63(39):849.
- SAMHSA. (2018). Risk and Protective Factors. Retrieved February 13, 2019, from <https://www.samhsa.gov/capt/practicing-effective-prevention/prevention-behavioral-health/risk-protective-factors>
- Seal, K. H., Kral, A. H., Gee, L., Moore, L. D., Bluthenthal, R. N., Lorvick, J., & Edlin, B. R. (2001). Predictors and prevention of nonfatal overdose among street-recruited injection heroin users in the San Francisco Bay Area, 1998-1999. American Journal of Public Health, 91(11), 1842–1846.
- Seal, Karen H., Thawley, R., Gee, L., Bamberger, J., Kral, A. H., Ciccarone, D., ... Edlin, B. R. (2005). Naloxone distribution and cardiopulmonary resuscitation training for injection drug users to prevent heroin overdose death: A pilot intervention study. Journal of Urban Health, 82(2), 303–311. <https://doi.org/10.1093/jurban/jti053>
- Tobin, K. E., Davey, M. A., & Latkin, C. A. (2005). Calling emergency medical services during drug overdose: an examination of individual, social and setting correlates. Addiction (Abingdon, England), 100(3), 397–404. <https://doi.org/10.1111/j.1360-0443.2005.00975.x>
- Walley, A. Y., Xuan, Z., Hackman, H. H., Quinn, E., Doe-Simkins, M., Sorensen-Alawad, A., ... Ozonoff, A. (2013). Opioid overdose rates and implementation of overdose education and nasal naloxone distribution in Massachusetts: interrupted time series analysis. BMJ, 346, f174. <https://doi.org/10.1136/bmj.f174>



USU EEO/Civil Rights Statement

In its programs and activities, including in admissions and employment, Utah State University does not discriminate or tolerate [discrimination](#), including harassment, based on race, color, religion, sex, national origin, age, genetic information, sexual orientation, gender identity or expression, disability, status as a protected veteran, or any other status protected by University policy, Title IX, or any other federal, state, or local law. Utah State University is an equal opportunity employer and does not discriminate or tolerate discrimination including harassment in employment including in hiring, promotion, transfer, or termination based on race, color, religion, sex, national origin, age, genetic information, sexual orientation, gender identity or expression, disability, status as a protected veteran, or any other status protected by University policy or any other federal, state, or local law. Utah State University does not discriminate in its housing offerings and will treat all persons fairly and equally without regard to race, color, religion, sex, familial status, disability, national origin, source of income, sexual orientation, or gender identity. Additionally, the University endeavors to provide reasonable accommodations when necessary and to ensure equal access to qualified persons with disabilities. The following office has been designated to handle inquiries regarding the application of Title IX and its implementing regulations and/or USU's non-discrimination policies: The Office of Equity in Distance Education, Room 400, Logan, Utah, titleix@usu.edu, 435-797-1266. For further information regarding non-discrimination, please visit equity.usu.edu, or contact: U.S. Department of Education, Office of Assistant Secretary for Civil Rights, 800-421-3481, ocr@ed.gov or U.S. Department of Education, Denver Regional Office, 303-844-5695 ocr.denver@ed.gov. Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Kenneth L. White, Vice President for Extension and Agriculture, Utah State University.

