

# Sage-grouse Brood Detection Techniques



**DAVID DAHLGREN  
TERRY MESSMER  
MIKE GUTTERY  
ERIC THACKER**

**UTAH STATE UNIVERSITY**

# Brood Detection Methods

- Walking
- Spotlighting
- Pointing Dogs



Photo  
by Les  
Flake

# Past Research



- Schroeder 1997
- Aldridge and Brigham 2001



Photo  
by Les  
Flake

“My dog, by the way, thinks that I have much to learn about partridges”

A. Leopold



Notice the  
breed of  
dog the  
“Father of  
Wildlife  
Science”  
chose!!!!

# Detection Methods



- Study Area
- Methods

Broods: N=21




Photo by Todd Black

# Spotlight



Walker et al. 2006 (WAFWA, Rapid City, South Dakota)

# Walking and Bird-dog Counts



Garmin™ Astro

# 2 Analyses



■ All Chicks

● Marked Only

○ Absolute Detection



# All Chick Analysis



- Overall difference ( $F = 7.89$ ,  $P = 0.001$ )


## Pair-wise comparisons

Walking differed from Bird-dog and Spotlight  
( $t = 3.85$ ,  $P = 0.001$  and  $t = 2.90$ ,  $P = 0.02$ )

Bird-dog flush and Spotlight counts did not differ  
( $t = 1.03$ ,  $P = 0.56$ )



# Marked Only Analysis



# Management Implications




Photo by  
Les Flake

Notice the Chicks



# Acknowledgments



- Technicians
- Grad students
- Committee
- PARM
- Jack H. Berryman Institute
- Other Funding Sources
  - UCWF
  - UDWR
  - USFS
  - NRCS
  - SITLA
  - Farm Bureau